

The Evening Sky Map

FREE* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

WWW.SKYPAPERS.COM

EQUATORIAL EDITION
NOVEMBER 2009

Sky Calendar – November 2009


- Mars 0.23° NNE from center of Beehive cluster (M44) (92° from Sun, morning sky) at 15h UT. Mag. +0.4.
- Full Moon at 19:14 UT.
- Venus 3.5° NNE from Spica (17° from Sun, morning sky) at 5h UT. Mags. -3.9 and +1.0.
- Moon near the Pleiades (morning sky) at 5h UT.
- Taurid (south) meteor shower peaks. Active between 25 Sept and 25 Nov. Associated with Comet 2P/Encke.
- Mercury at superior conjunction with the Sun at 8h UT. The planet passes into the evening sky.
- Moon at perigee (closest to Earth) at 7h UT (368,903 km; 32.9').
- Moon near Beehive cluster (M44) (99° from Sun, morning sky) at 23h UT.
- Moon near Mars (morning sky) at 14h UT. Mag. +0.3.
- Last Quarter Moon at 15:56 UT.
- Moon near Regulus (morning sky) at 13h UT.
- Taurid (north) meteor shower peaks. May produce the occasional bright fireball.
- Moon near Saturn (morning sky) at 20h UT. Mag. +1.1.
- Moon near Spica (morning sky) at 12h UT.
- New Moon at 19:13 UT. Start of lunation 1075.
- Leonid meteor shower peaks at 9h UT. Arises from debris ejected by Comet Tempel-Tuttle in 1533. Expect about 25 to 30 meteors per hour under dark skies. Predictions of enhanced activity between 21–22h UT on 17 Nov (favours sky watchers in Asia).
- Alpha Monocerotid meteor shower peaks at 15:25 UT. A usually minor shower active 15–25 Nov. Radiant is near Procyon. Predictions of enhanced activity this year. Timing favours Far East Asia, Australia and across the Pacific to Alaska.
- Moon at apogee (farthest from Earth) at 20h UT (distance 404,733 km; angular size 29.7').
- Moon near Jupiter (evening sky) at 19h UT. Mag. -2.3.
- First Quarter Moon at 21:38 UT.

More sky events and links at <http://Skymaps.com/skycalendar/>
All times in Universal Time (UT). (Singapore Standard Time = UT + 8 hours.)

SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY NOV 9 PM
LATE NOV 8 PM

SKY MAP DRAWN FOR A LATITUDE OF 0° AND IS SUITABLE FOR LATITUDES UP TO 15° NORTH OR SOUTH OF THIS



Symbols

- Galaxy
- Double Star
- Variable Star
- Diffuse Nebula
- Planetary Nebula
- Open Star Cluster
- Globular Star Cluster

Star Magnitudes -1 0 1 2 3 4

Sky maps Follow us on Twitter
<http://twitter.com/skymaps>

International Year of Astronomy 2009
<http://www.astronomy2009.org>

Copyright © 2000–2009 Kym Thalassoudis. All Rights Reserved.
* TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. Singapore Standard Time is UT plus 8 hours.

Variable Star – A star that changes brightness over a period of time.

EQUATORIAL EDITION
NOVEMBER 2009

CELESTIAL OBJECTS

☆
☆
☆
☆
Sky maps .com

Easily Seen with the Naked Eye

Altair	Aql	●	Brightest star in Aquila. Name means "the flying eagle". Dist=16.8 ly.
Capella	Aur	●	The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.
δ Cephei	Cep	☉	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg	●	Brightest star in Cygnus. One of the greatest known supergiants. Dist=1,400±200 ly.
Achernar	Eri	●	Brightest star in Eridanus, The River. Arabic name meaning "end of river". Dist=144 ly.
Vega	Lyr	●	The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly.
Rigel	Ori	●	The brightest star in Orion. Blue supergiant star with mag 7 companion. Dist=770 ly.
Betelgeuse	Ori	●	One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Algol	Per	☉	Famous eclipsing binary star. Magnitude varies between 2.1 & 3.4 over 2.867 days.
Fomalhaut	PsA	●	Brightest star in Piscis Austrinus. In Arabic the "fish's mouth". Dist=25 ly.
Pleiades	Tau	☉	The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=399 ly.
Hyades	Tau	☉	Large V-shaped star cluster. Binoculars reveal many more stars. Dist=152 ly.
Aldebaran	Tau	●	Brightest star in Taurus. It is not associated with the Hyades star cluster. Dist=66.7 ly.
Polaris	UMi	●	The North Pole Star. A telescope reveals an unrelated mag 8 companion star. Dist=433 ly.

Easily Seen with Binoculars

M31	And	☉	The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.5 million ly.
M2	Aqr	☉	Resembles a fuzzy star in binoculars.
η Aquilae	Aql	☉	Bright Cepheid variable. Mag varies between 3.6 & 4.5 over 7.166 days. Dist=1,200 ly.
M38	Aur	☉	Stars appear arranged in "pi" or cross shape. Dist=4,300 ly.
M36	Aur	☉	About half size of M38. Located in rich Milky Way star field. Dist=4,100 ly.
μ Cephei	Cep	☉	Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
Mira	Cet	☉	Famous long period variable star. Mag varies between 3.0 & 10.1 over 332 days.
χ Cygni	Cyg	☉	Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
M39	Cyg	☉	May be visible to the naked eye under good conditions. Dist=900 ly.
LMC	Dor	☉	Large Magellanic Cloud. A neighbouring galaxy of the Milky Way. Dist=180,000 ly.
ε Lyrae	Lyr	●	Famous Double Double. Binoculars show a double star. High power reveals each a double.
R Lyrae	Lyr	☉	Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
Cr 69	Ori	☉	Lambda Orionis Cluster. Dist=1,630 ly.
M42	Ori	☐	The Great Orion Nebula. Spectacular bright nebula. Best in telescope. Dist=1,300 light years.
κ Pavonis	Pav	☉	Cepheid-type. Magnitude varies between 3.9 & 4.8 over 9.088 days.
6752	Pav	☉	One of the better globular star clusters in the sky. Dist=14,000 ly.
M15	Peg	☉	Only globular known to contain a planetary nebula (Mag 14, d=1"). Dist=30,000 ly.
Double Cluster	Per	☉	Double Cluster in Perseus. NGC 869 & 884. Excellent in binoculars. Dist=7,300 ly.
ζ Phoenicis	Phe	☉	Eclipsing binary star and double (mag 8). Varies between 3.9 & 4.4 over 1.667 days.
253	Scl	☉	Fine, large, cigar-shaped galaxy. Requires dark sky. Member of Sculptor Group.
47 Tucanae	Tuc	☉	Spectacular object. Telescope will reveal stars. Near edge of SMC. Dist=15,000 ly.
β Tucanae	Tuc	●	Complex multiple star. Binoculars show one pair. Telescope required to split primary star.
SMC	Tuc	☉	Small Magellanic Cloud. Companion galaxy to Milky Way. Requires dark sky. Dist=210,000 ly.
Cr 399	Vul	☉	Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.

Telescopic Objects

γ Andromedae	And	●	Attractive double star. Bright orange star with mag 5 blue companion. Sep=9.8".
7009	Aqr	✦	Saturn Nebula. Requires 8-inch telescope to see Saturn-like appendages.
7293	Aqr	✦	Helix Nebula. Spans nearly 1/4 deg. Requires dark sky. Dist=300 ly.
γ Arietis	Ari	●	Impressive looking double blue-white star. Visible in a small telescope. Sep=7.8".
η Cassiopeiae	Cas	●	Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".
Albireo	Cyg	●	Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".
61 Cygni	Cyg	●	Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".
γ Delphini	Del	●	Appear yellow & white. Mags 4.3 & 5.2. Dist=100 ly. Struve 2725 double in same field.
θ Eridani	Eri	●	Striking blue-white double star. Mags 3.2 & 4.3. Visible in a small telescope. Sep=8.2".
β Lyrae	Lyr	☉	Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star.
M57	Lyr	✦	Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.
M11	Sct	☉	Wild Duck Cluster. Resembles a globular through binoculars. V-shaped. Dist=5,600 ly.
M1	Tau	☐	Crab Nebula. Remnant from supernova which was visible in 1054. Dist=6,500 ly.
M33	Tri	☉	Fine face-on spiral galaxy. Requires a large aperture telescope. Dist=2.3 million ly.
M27	Vul	✦	Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.