

# The Evening Sky Map

FREE\* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

## Sky Calendar – April 2019

Get Sky Calendar on Twitter  
<http://twitter.com/skymaps>

- Moon at apogee** (farthest from Earth) at 0h UT (distance 405,577 km; angular size 29.5').
- Moon near Venus** (morning sky) at 7h UT. Mag. -4.0.
- Moon near Mercury** (26° from Sun, morning sky) at 2h UT. Mag. 0.8.
- New Moon** at 8:51 UT. Start of lunation 1191.
- Moon near the Pleiades** (evening sky) at 23h UT.
- Moon near Mars** (evening sky) at 10h UT. Mag. 1.5.
- Moon near Aldebaran** (evening sky) at 16h UT.
- Mercury at greatest elongation west** (28° from Sun, morning sky) at 20h UT. Mag. 0.4.
- First Quarter Moon** at 19:05 UT.
- Moon near Pollux** (evening sky) at 22h UT.
- Moon near Beehive cluster M44** (evening sky) at 21h UT.
- Mars 6.5° N of Aldebaran** (45° from Sun, evening sky) at 1h UT. Mags. 1.6 and 0.9.
- Moon near Regulus** (evening sky) at 10h UT.
- Mercury 4.3° E of Venus** (30° from Sun, morning sky) at 20h UT. Mags. 0.2 and -3.9.
- Moon at perigee** (closest to Earth) at 22:03 UT (364,205 km; angular size 32.8').
- Moon near Spica** (morning sky) at 3h UT.
- Full Moon** at 11:11 UT.
- Moon near Antares** (morning sky) at 11h UT.
- Lyrid meteor shower** peaks at 0h UT. Active April 14-30. Radiant is between Hercules and Lyra. Expect 10 to 20 bright, fast meteors per hour at its peak. Unfortunately, bright moonlight this year means poor viewing conditions.
- Moon near Jupiter** (morning sky) at 13h UT. Mag. -2.4.
- Last Quarter Moon** at 22:18 UT.
- Moon near Saturn** (105° from Sun, morning sky) at 13h UT. Mag. 0.5. Occultation visible from eastern Australia, New Zealand, and western South America.
- Moon at apogee** (farthest from Earth) at 18h UT (distance 404,582 km; angular size 29.5').


More sky events and links at <http://Skymaps.com/skycalendar/>

All times in Universal Time (UT). (Singapore Standard Time = UT + 8 hours.)



SAVE ON RECOMMENDED PRODUCTS • <http://Skymaps.com/store>

- STAR ATLASES & PLANISPHERES
  - STAR CHARTS & ASTRO POSTERS
  - BOOKS FOR SKY WATCHERS
  - TELESCOPES & BINOCULARS
- All sales support the production and free distribution of The Evening Sky Map.



EQUATORIAL EDITION  
 APRIL 2019

SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY APR 8 PM  
 LATE APR 7 PM

SKY MAP DRAWN FOR A LATITUDE OF 0° AND IS SUITABLE FOR LATITUDES UP TO 15° NORTH OR SOUTH OF THIS

### Symbols

- Galaxy ☾
- Double Star ●●
- Variable Star ⊙
- Diffuse Nebula □
- Planetary Nebula ☆
- Open Star Cluster ○
- Global Star Cluster ⊕

Star Magnitudes ●●●●●  
 -1 0 1 2 3 4

Copyright © 2000–2019 Kym Thalassoudis. All Rights Reserved.

\*TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

INSTRUCTIONS: THE SKY MAP SHOWS THE ENTIRE NIGHT SKY FROM HORIZON-TO-HORIZON AS IT APPEARS ON CERTAIN DATES AND TIMES. THE CENTER OF THE MAP IS THE PART OF THE SKY DIRECTLY OVERHEAD (ZENITH) AND THE OUTER RING IS THE PART OF THE SKY DIRECTLY ON THE HORIZON. CELESTIAL OBJECTS ARE LOCATED BETWEEN THE ZENITH AND THE HORIZON. COMPASS DIRECTIONS ARE INDICATED ALONG THE HORIZON CIRCLE (FOR EXAMPLE, "NORTH"). TURN THE SKY MAP AROUND ITS CENTER TO FIND A BRIGHT STAR PATTERN IN THE SKY.

## About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

## Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

## Astronomical Glossary

**Conjunction** – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

**Constellation** – A defined area of the sky containing a star pattern.

**Diffuse Nebula** – A cloud of gas illuminated by nearby stars.

**Double Star** – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

**Ecliptic** – The path of the Sun's center on the celestial sphere as seen from Earth.

**Elongation** – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

**Galaxy** – A mass of up to several billion stars held together by gravity.

**Globular Star Cluster** – A ball-shaped group of several thousand old stars.

**Light Year (ly)** – The distance a beam of light travels at 300,000 km/sec in one year.

**Magnitude** – The brightness of a celestial object as it appears in the sky.

**Open Star Cluster** – A group of tens or hundreds of relatively young stars.

**Opposition** – When a celestial body is opposite the Sun in the sky.

**Planetary Nebula** – The remnants of a shell of gas blown off by a star.

**Universal Time (UT)** – A time system used by astronomers. Also known as Greenwich Mean Time. Singapore Standard Time is UT plus 8 hours.

**Variable Star** – A star that changes brightness over a period of time.

EQUATORIAL EDITION  
APRIL 2019

CELESTIAL OBJECTS

★  
★  
★  
★  
Sky maps .com

## Easily Seen with the Naked Eye

Sirius	CMa	•	The brightest star in the sky. Also known as the "Dog Star". Dist=8.6 ly.
Procyon	CMi	•	Greek name meaning "before the dog" -- rises before Sirius (northern latitudes). Dist=11.4 ly.
Canopus	Car	•	Second brightest star in the sky. 14,000 times more luminous than the Sun. Dist=310 ly.
Castor	Gem	•	Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	•	With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
Regulus	Leo	•	Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Betelgeuse	Ori	•	One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Pleiades	Tau	☉	The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=380 ly.
Hyades	Tau	☉	Large V-shaped star cluster. Binoculars reveal many more stars. Dist=151 ly.
Spica	Vir	•	Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=260 ly.

## Easily Seen with Binoculars

M44	Cnc	☉	Praesepe or Beehive Cluster. Visible to the naked eye. Dist=577 ly.
M41	CMa	☉	First recorded observation by Aristotle in 325 BC as "cloudy spot". Dist=2,300 ly.
2516	Car	☉	Spectacular open star cluster of 100 stars spanning 1/2 deg. Dist=1,300 ly.
2808	Car	☉	Located 4 deg W of Nu Carinae. Visible to the naked eye on clear nights.
R Carinae	Car	☉	Long period variable. Magnitude varies between 3.9 & 10.5 over 309 days.
3114	Car	☉	Stunning open cluster. 30+ stars visible through 7x binoculars. Dist=2,900 ly.
3293	Car	☉	Rich, tightly packed. Surrounded by large, faint nebulosity. Dist=8,500 ly.
IC 2602	Car	☉	The Southern Pleiades. Bright cluster twice diameter of full Moon. Dist=500 ly.
3372	Car	☐	Eta Carinae Nebula. Enormous glowing cloud in rich star field. Dist=8,000 ly.
3532	Car	☉	Herschel -- "most brilliant cluster". 60+ stars in 7x binoculars. Dist=1,300 ly.
ω Centauri	Cen	☉	Largest and brightest globular star cluster in sky. 1 million stars. Dist=17,000 ly.
Mel 111	Com	☉	Coma Berenices. 80 mag 5-6 stars in 5 deg. Dist=260 ly. Age=400 million years.
LMC	Dor	☉	Large Magellanic Cloud. A neighbouring galaxy of the Milky Way. Dist=180,000 ly.
M35	Gem	☉	Fine open cluster located near foot of the twin Castor. Dist=2,800 ly.
M48	Hya	☉	12+ stars in 7x binoculars. Triangular asterism near centre. Dist=1,990 ly.
γ Leporis	Lep	•	Visible with binoculars. Gold & white stars. Mags 3.6 & 6.2. Dist=30 ly. Sep=96.3".
2232	Mon	☉	A large scattered star cluster of 20 stars. Dist=1,300 ly.
2244	Mon	☉	Surrounded by the rather faint Rosette Nebula. Dist=5,540 ly.
M50	Mon	☉	Visible with binoculars. Telescope reveals individual stars. Dist=3,000 ly.
M42	Ori	☐	The Great Orion Nebula. Spectacular bright nebula. Best with telescope. Dist=1,300 light years.
L <sup>2</sup>	Pup	☉	Semi-regular variable. Magnitude varies between 2.6 & 6.2 over 140.42 days.
M47	Pup	☉	Bright star cluster. 15+ stars in 7x binoculars. Dist=1,500 ly.
M46	Pup	☉	Dist=5,400 ly. Contains planetary NGC 2438 (Mag 11, d=65") -- not associated.
2451	Pup	☉	30+ stars in binoculars. The brightest star, χ Puppis, is red. Dist=850 ly.
2477	Pup	☉	Very rich but distant star cluster (4,200 ly). Resembles globular through binoculars.
2547	Vel	☉	Fine open cluster visible through binoculars. Dist=1,300 ly.
IC 2391	Vel	☉	Omicron Velorum Cluster. Superb object for binoculars. Dist=450 ly.

## Telescopic Objects

M67	Cnc	☉	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.
M94	CVn	☉	Compact nearly face-on spiral galaxy. Dist=15 million ly.
M64	Com	☉	Black-Eye Galaxy. Discovered by J.E. Bode in 1775 -- "a small, nebulous star".
2070	Dor	☐	Tarantula Nebula. A bright nebula located in LMC. A star-forming region.
3242	Hya	☉	Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.
M83	Hya	☉	Classic face-on spiral. Discovered in 1752 by Lacaille. In attractive star field.
γ Leonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".
β Monocerotis	Mon	•	Triple star. Mags 4.6, 5.0 & 5.4. Requires telescope to view arc-shape. Sep=7.3".
k Puppis	Mon	•	Christmas Tree Cluster. Associated with the Cone Nebula. Dist=2,450 ly.
M1	Pup	•	Telescope easily shows two blue-white stars of almost equal brightness. Sep=9.9".
M81	Tau	☐	Crab Nebula. Remnant from supernova which was visible in 1054. Dist=6,500 ly.
M82	UMa	☉	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.
M82	UMa	☉	Close to M81 but much fainter and smaller.
3132	Vel	☉	One of the brightest planetaries. Magnitude 10 central star. Dist=2,600 ly.
M104	Vir	☉	Sombrero Galaxy. Almost edge-on spiral galaxy. Protruding central core.
γ Virginis	Vir	•	Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.